



GROUP M FUNGICIDE

Daconil[®] 2787 Flowable Fungicide

COMMERCIAL

Turf and Ornamental Fungicide

GUARANTEE:

Chlorothalonil (tetrachloroisophthalonitrile)..... 40.4%
Contains 500 grams Chlorothalonil per Liter

**READ THE LABEL AND PAMPHLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN**

WARNING: CAUSES SEVERE EYE DAMAGE
See the side panel for additional precautionary statements

REGISTRATION NUMBER: **15724**
PEST CONTROL PRODUCTS ACT

NET CONTENTS : **10L**

Syngenta Crop Protection Canada, Inc.
140 Research Lane, Research Park
Guelph, Ontario
N1G 4Z3

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offence under the *Pest Control Products Act* to use a control product under unsafe conditions.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed, IMMEDIATELY contact Syngenta's Emergency Information Network at 1-800-F-A-S-T-M-E-D (327-8633). Do not induce vomiting. Vomiting should only be induced under the direction of a physician or a poison control center. If spontaneous vomiting occurs, have victim lean forward with head down to avoid breathing in of vomitus, rinse mouth and administer water. Never give anything by mouth to an unconscious person. Obtain medical attention or contact a poison control centre **IMMEDIATELY**.

In case of contact with eyes, IMMEDIATELY flush eyes with plenty of water and continue for 15 minutes. Seek medical attention for eyes **IMMEDIATELY**. Persons having allergic reaction should contact a physician. Affected persons respond to treatment with antihistamines or steroid creams and/or systemic steroids.

In case of contact with skin, flush skin with running water for a minimum of 20 minutes. Start flushing while removing contaminated clothing. If irritation persists, repeat flushing. Obtain medical attention **IMMEDIATELY**.

If inhaled, remove victim to fresh air. Until adequate ventilation or other control measures are restored, employ proper respiratory protection. If symptoms develop, seek medical attention.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. Warning: Causes Severe Eye Damage.

DO NOT get in eyes. Wear protective goggles or eye shield when handling this product. AVOID contact with skin or clothing. Wash exposed areas of skin with soap and warm water after handling or using. DO NOT take internally. AVOID breathing spray mist. This product may produce temporary allergic side effects, characterized by redness of the eyes, mild bronchial irritation and a redness or rash on exposed skin areas. DO NOT store near feed or foodstuffs. Store in a cool place. Protect from excessive heat. Apply only to areas specified on label.

This product is toxic to fish, aquatic invertebrates and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. DO NOT apply directly to water.

Keep out of lakes, streams, and ponds. DO NOT contaminate water by cleaning of equipment, or disposal of wastes. DO NOT apply where runoff is likely to occur. DO NOT apply when weather conditions favour drift from areas being treated.

TOXICOLOGICAL INFORMATION

Treat symptomatically

DISPOSAL OF UNUSED, UNWANTED PRODUCT

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean up of spills.

CONTAINER DISPOSAL:

For recyclable containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING
CALL 1-800-327-8633 (FASTMED)***



| | | |
|-------|----------|-----------|
| GROUP | M | FUNGICIDE |
|-------|----------|-----------|

Daconil[®] 2787 Flowable Fungicide

COMMERCIAL

Turf and Ornamental Fungicide

GUARANTEE:

| | |
|--|-------|
| Chlorothalonil (tetrachloroisophthalonitrile)..... | 40.4% |
| Contains 500 grams Chlorothalonil per Liter | |

READ THE LABEL AND PAMPHLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

WARNING: CAUSES SEVERE EYE DAMAGE
 See the side panel for additional precautionary statements

REGISTRATION NUMBER: **15724**
 PEST CONTROL PRODUCTS ACT

Syngenta Crop Protection Canada, Inc.
 140 Research Lane, Research Park
 Guelph, Ontario
 N1G 4Z3

NOTICE TO USER

This control product is to be used only in accordance with the directions on this label. It is an offence under the *Pest Control Products Act* to use a control product under unsafe conditions.

NOTICE TO BUYER

Seller's guarantee shall be limited to the terms set out on the label and, subject thereto, the buyer assumes the risk to persons or property arising from the use or handling of this product and accepts the product on that condition.

FIRST AID

IN CASE OF POISONING, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed, IMMEDIATELY contact Syngenta's Emergency Information Network at 1-800-F-A-S-T-M-E-D (327-8633). Do not induce vomiting. Vomiting should only be induced under the direction of a physician or a poison control center. If spontaneous vomiting occurs, have victim lean forward with head down to avoid breathing in of vomitus, rinse mouth and administer water. Never give anything by mouth to an unconscious person. Obtain medical attention or contact a poison control centre **IMMEDIATELY**.

In case of contact with eyes, IMMEDIATELY flush eyes with plenty of water and continue for 15 minutes. Seek medical attention for eyes **IMMEDIATELY**. Persons having allergic reaction should contact a physician. Affected persons respond to treatment with antihistamines or steroid creams and/or systemic steroids.

In case of contact with skin, flush skin with running water for a minimum of 20 minutes. Start flushing while removing contaminated clothing. If irritation persists, repeat flushing. Obtain medical attention **IMMEDIATELY**.

If inhaled, remove victim to fresh air. Until adequate ventilation or other control measures are restored, employ proper respiratory protection. If symptoms develop, seek medical attention.

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN. Warning: Causes Severe Eye Damage.

DO NOT get in eyes. Wear protective goggles or eye shield when handling this product. AVOID contact with skin or clothing. Wash exposed areas of skin with soap and warm water after handling or using. DO NOT take internally. AVOID breathing spray mist. This product may produce temporary allergic side effects, characterized by redness of the eyes, mild bronchial irritation and a redness or rash on exposed skin areas. DO NOT store near feed or foodstuffs. Store in a cool place. Protect from excessive heat. Apply only to areas specified on label.

This product is toxic to fish, aquatic invertebrates and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. DO NOT apply directly to water.

Keep out of lakes, streams, and ponds. DO NOT contaminate water by cleaning of equipment, or disposal of wastes. DO NOT apply where runoff is likely to occur. DO NOT apply when weather conditions favour drift from areas being treated.

TOXICOLOGICAL INFORMATION

Treat symptomatically

DISPOSAL OF UNUSED, UNWANTED PRODUCT

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean up of spills.

CONTAINER DISPOSAL:

For recyclable containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING
CALL 1-800-327-8633 (FASTMED)***

PRODUCT INFORMATION

A BROAD SPECTRUM AGRICULTURAL FUNGICIDE FOR USE ON GOLF COURSE TEES, GREENS AND FAIRWAYS, ORNAMENTAL TURFGRASS AND ORNAMENTALS.

Read entire label carefully and use only as directed.

DACONIL 2787 Flowable Fungicide is formulated for use on golf course tees, greens and fairways, ornamental turfgrass and ornamental herbs, shrubs and trees. It is highly effective for the control of a broad spectrum of turf and ornamental plant diseases when it is used according to the directions on this product label. Thorough, uniform coverage of plant surfaces is essential for good disease control.

Mixing Instructions

DACONIL 2787 Flowable Fungicide Only

Apply DACONIL 2787 Flowable Fungicide as a spray, using sufficient water to obtain adequate coverage of plant surfaces. Volume of spray to be used may vary, depending on crop and plant growth, from 200 to 4,000 liters per hectare for dilute sprays, and 50 to 100 liters per hectare for concentrate sprays. Use the higher spray volumes when foliage is dense and coverage is more difficult.

Note: Before preparing the spray mix, slowly invert the DACONIL 2787 Flowable Fungicide container several times to assure uniform mixture.

Add the required amount of DACONIL 2787 Flowable Fungicide to the spray tank while filling with water. Keep agitator running when filling spray tank and during spray operations. DO NOT combine DACONIL 2787 Flowable Fungicide in the spray tank with pesticides, surfactants or fertilizers unless your prior use has shown the combination physically compatible, effective and non injurious under your conditions of use.

DO NOT APPLY BY AIR

DIRECTIONS FOR USE

Sod farm turf treated with Chlorothalonil prior to harvest must be mechanically cut, rolled and harvested.

This product is not to be used around homes or other residential areas such as parks, school grounds, playing fields. It is not for use by homeowners or other uncertified users.

Do not apply this product in a way that will contact workers or other persons or pets either directly or through drift. Only protected handlers may be in the area during application.

Turf: DO NOT mow or water after treatment until spray deposit on turfgrass is thoroughly dry; DACONIL 2787 Flowable Fungicide should always be used in conjunction with good turf management practices.

Golf Course Fairways: Apply DACONIL 2787 Flowable Fungicide when conditions favor disease development and repeat applications as long as these conditions persist. Under severe disease conditions use the highest rate and shortest interval corresponding with the application schedule selected from the table below.

| Disease | Application Interval | Application Rate Litres Per Hectare |
|---------------------------|-----------------------------|--|
| Sclerotinia dollar spot | 7-14 days | 4.8 to 9.5 |
| Helminthosporium leafspot | 7-14 days | 9.5 to 19.0 |
| Rhizoctonia brown patch | 7-14 days | 9.5 to 19.0 |

Golf Course Tees and Greens and Ornamental Turfgrass: Apply DACONIL 2787 Flowable Fungicide in an adequate amount of water to provide complete coverage. This amount may vary from 8 to 40 liters per 100 sq. metres. See below for suggested rates and timing. Under severe disease conditions, use the curative rates and spray on a 7 day schedule.

| Disease | Interval of Application | RATE: Millilitres per 100 sq. Metres | |
|---|-------------------------|--------------------------------------|-----------------|
| | | Preventive* | Curative** |
| Dollar spot | 7-14 days | 95-190**** | 190-350 |
| Helminthosporium leafspot and melting out | 7-10 days | 95-190 | 190-350 |
| Large brown patch | 7-10 days | 95-190 | 190-350 |
| Anthrachnose | 7-10 days | 170-190 | not-recommended |
| Gray snow mould (<i>Typhula incarnata</i>) and Pink snow mould (<i>Microdochium nivale</i>) | – | 240-500*** | not-recommended |

* Recommended rates for preventing disease establishment; use lower rate as when disease conditions are light to moderate, higher indicated rates when conditions are severe.

** Rates for use on a 7day schedule when disease is present. Higher indicated rate should be applied under severe conditions.

*** Apply DACONIL 2787 Flowable Fungicide at 500 milliliters or apply 240 milliliters as a tank mix with 125 to 250 milliliters of ROVRAL® Green 240 g/L per 100 sq. metres prior to snow cover in the fall and repeat in the early spring if conditions for snow mould continue to exist.

**** Apply DACONIL 2787 Flowable Fungicide at 190 milliliters as a tank mix with 26 milliliters of Banner MAXX® Fungicide per 100sq. metres and an interval of 21-28 days to control Dollar Spot.

Tank Mix Instructions:

DACONIL 2787 Flowable Fungicide can be tank mixed with Primo® MAXX Turf Growth Regulator or Rovral® Green for control of Grey snow mould and Pink snow mould only. Read the Primo MAXX, Banner MAXX and Rovral Green labels and follow the precautionary statements and directions for use. Follow the most restrictive instructions and the greatest interval of application on the labels of the products to be tank-mixed.

1. Always check the compatibility of Primo MAXX, Banner MAXX or Rovral Green tank mix using a jar test with proportionate amounts of DACONIL 2787 Flowable Fungicide, and the water, before mixing in the spray tank.
2. Provide sufficient jet or mechanical agitation during filling and application to keep the tank mix uniformly suspended.
3. Fill tank at least 1/2 full of clean water.
4. If using Rovral Green, Primo MAXX or Banner MAXX as a tank mix partner add them first then add DACONIL 2787 Flowable Fungicide next.
5. Do not leave tank mix combinations in the spray tank for prolonged periods without agitation. Mix and apply them the same day.

Ornamentals and Conifers: Apply DACONIL 2787 Flowable Fungicide at rate of 2.5 liters per 1000 liters of water unless other directions are given in the tables below. For ornamentals, apply sufficient spray to obtain adequate coverage, but do not exceed 2000 liters of water per hectare. Begin applications as directed for each species and disease condition cited and repeat on a 7 to 14 day schedule until conditions are no longer favorable for disease development. During periods when conditions favor severe disease incidence, generally cloudy or wet weather, use the higher rate specified and the shortest indicated interval between applications.

DACONIL 2787 Flowable Fungicide may be used in greenhouse. Applicators and attending personnel should wear protective clothing including long sleeves, gloves, goggles or eye shield plus a face-fitting respirator specifically designed to remove organic pesticide vapors and particulates. DO NOT use mistblowers or high pressure spray equipment when making applications of DACONIL 2787 Flowable Fungicide in greenhouses. Use of DACONIL 2787 Flowable Fungicide is recommended for disease control on the ornamentals and conifers listed in the tables below. Use of

DACONIL 2787 Flowable Fungicide is not recommended for species that are not listed.

ORNAMENTALS:

| Species | Diseases Controlled | Suggested First Application |
|------------------------------------|---|-----------------------------------|
| BROADLEAF SHRUBS AND TREES: | | |
| Ash (Fraxinus) | Cercospora, Cercosporidium, Cylindrosporium leafspots | Spring bud break |
| Azalea, *Rhododendron* | Phytophthora die-back; Ovulinia flower blight | New leaf emergence Early bloom |
| Buckeye, Horsechestnut | Leaf blotch, Anthracnose | Spring bud break |
| Cherry B Laurel | Cercospora leafspot | Petal fall |
| Crabapple | Scab, Cedar-apple rust, Sphaeropsis leafspot | Spring bud break |
| Dogwood | Septoria leafspot | Early bloom |
| Euonymus | Anthracnose | Spring bud break |
| Firethorn | Scab | Spring bud break |

| Species | Diseases Controlled | Suggested First Application |
|---------------------------------------|---|-----------------------------|
| BROADLEAF SHRUBS AND TREES: | | |
| Flowering Almond, Quince, Sand Cherry | Monilinia blossom/branch blight | Early bloom |
| Hawthorn | Rust, Fabraea leafspot | Pre-bloom |
| Holly | Rhizoctonia web blight | Warm, moist conditions |
| Mountain Laurel | Cercospora leafspot | Spring bud break |
| Oak (red group only) | Taphrina blister, Actinopelte leafspot, Anthracnose | Dormant budswell |
| Oregon B Grape (Mahonia) | Rust | Spring bud break |
| Photinia | Fabraea (Entomosporium) leafspot | Spring bud break |
| Pieris (Andromeda) | Phytophthora die-back | New leaf emergence |
| Poplar | Marssonina leafspot | Spring bud break |
| Privet | Cercospora leafspot | Prolonged wet conditions |
| Sycamore, Planetree | Anthracnose | Spring bud break |
| Viburnum | Powdery mildew | Mid-summer |

| Species | Diseases Controlled | Suggested First Application |
|---|---|--|
| BULBS AND FLOWERING PLANTS: | | |
| Carnation | Alternaria leafspot/branch rot: Botrytis flower blight | Transplant of cuttings; cool, moist conditions |
| Chrysanthemum/Daisy | Mycosphaerella ray blight, Septoria leafspot; Botrytis flower blight (gray mould) | Transplant of cuttings Pre-bloom |
| Geranium | Botrytis blight, rust | Cool, moist conditions |
| Gladiolus | Curvularia leaf/ flower spot, Botrytis leaf/flower spot | Early propagation |
| Hollyhock | Rust | Early seedling stage |
| Hydrangea* (foliage only) | Cercospora and Septoria leafspots, Rust | Early propagation |
| Iris | Botrytis blossom blight, Didymellina leafspot | Cool, moist conditions |
| Lily | Botrytis gray mould | Pre-bloom |
| Petunia* | Phytophthora blight (foliar phase), Botrytis blight | Pre-bloom |
| Rose (Use 1.8 L per 1000 L) | Black spot, Botrytis blight | Spring bud break |
| Statice | Anthracnose, Cercospora, Alternaria, Botrytis leaf blights | Spring bud break |
| Zinnia | Powdery mildew | First sign of disease |
| * Discolouration of blooms has been noted on certain varieties when applications are made during flowering. | | |

| Species | Diseases Controlled | Suggested First Application |
|-------------------------------------|---|-----------------------------|
| FOLIAGE PLANTS | | |
| Dracaena | Fusarium leafspot | Pre-transplant |
| Pachysandra (Use 5 L per 1000 L) | Volutella leaf blight | Spring bud break |
| Leatherleaf fern | Ascochyta blight, Cercospora leafspot, Cylindrocladium leafspot, Rhizoctonia blight | Spring bud break |
| Parlor palm (Chamaedorea) | Bipolaris (Helminthosporium) leafspot | Cool, moist conditions |
| Prayer plant (Maranta) | Helminthosporium leafspot | Early propagation |
| Oyster plant (Rhoeo) | Tan leafspot | Early propagation |
| Syngonium | Cephalosporium leafspot | Warm, moist conditions |
| Philodendron | Phytophthora blight, Dactylaria leafspot | Moist conditions |

CONIFERS:

| Diseases Controlled | DACONIL 2787 Flowable Fungicide Rate/Hectare | Application Directions |
|--|---|--|
| Scleroderris Canker (pines), Swiss Needlecast and Rhabdocline Needlecast (Douglas fir) | 2.4 to 4.8 L | Apply 100 to 1000 liters of water per hectare. Make first application in spring when new shoot growth is 1 to 5 cm in length. Make additional applications 3 to 4 week intervals until conditions no longer favor disease development. For use in nursery beds, apply the highest rate specified on a 3 week schedule. |
| Sirococcus Tip Blight | 3.6 to 6 L | |
| Rhizosphaera Needlecast (spruces), Scirrhia brown spot (pines) | 9.5 L | |
| Lophodermium Needlecast | 2.4 to 4.8 L | Apply in 100 to 1000 liters of water per hectare. Begin applications in mid-July to early August before infection occurs. Make additional applications at 3 to 4 week intervals until conditions no longer favor disease development. For use in nursery beds, apply 4.8 L/ha on a 3 week schedule. |
| Botrytis seedling blight, Phoma twig blight | 2.4 to 4.8 L | Apply in 100 to 1000 liters of water per hectare. Begin applications in nursery beds when seedlings are 10 cm tall and when cool, moist conditions favor disease development. Make additional applications at 7 to 14 day intervals as long as disease favorable conditions persist. |

Resistance Management Recommendations

For resistance management, DACONIL 2787 Flowable Fungicide contains a Group M fungicide. Any fungal population may contain individuals naturally resistant to DACONIL 2787 Flowable Fungicide and other Group M fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To delay fungicide resistance:

- Where possible, rotate the use of DACONIL 2787 Flowable Fungicide or other Group M fungicides with different groups that control the same pathogens.
- Avoid application of more than the maximum number listed in the label and consecutive sprays of DACONIL 2787 Flowable Fungicide or other fungicides in the same group in a season.
- Fungicide use should be based on an IPM program that includes scouting, historical information related to pesticide use and crop rotation and considers cultural, biological and other chemical control practices.
- Monitor treated fungal populations for sign of resistance development. If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different target site of action, if available.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and disease problems in your area.

For further information or to report suspected resistance, contact company representatives at 1-800-459-2422 (in Eastern Canada) or 1-800-665-9250 (in Western Canada) or at www.syngenta.ca.

19-APR-2002

CORRECTIVE SUPPLEMENTAL LABEL

DACONIL® 2787 FLOWABLE FUNGICIDE

Tank-Mix Rate Correction with ROVRAL® Green 240 g/L

Turf and Ornamental Fungicide

GUARANTEE:

Chlorothalonil (tetrachloroisophthalomitrile).....40.4%

Contains 500 grams Chlorothalonil per Liter

READ THE LABEL AND BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

WARNING: Causes Severe Eye Damage

See the label for additional precautionary statements

REGISTRATION NUMBER: **15724**

PEST CONTROL PRODUCTS ACT

Syngenta Crop Protection Canada, Inc.

140 Research Lane

Research Park, University of Guelph

Guelph, Ontario

N1G 4Z3

CORRECTION TO DIRECTIONS FOR USE - Tank mix with ROVRAL Green 240 g/L

Golf Course Tees and Greens and Ornamental Turfgrass: Apply DACONIL 2787 in an adequate amount of water to provide complete coverage. This amount may vary from 8 to 40 liters per 100 sq. metres. See below for suggested rates and timing. Under severe disease conditions, use the curative rates and spray on a 7 day schedule.

| Disease | Interval of Application | RATE: Millilitres per 100 sq. Metres | |
|---|-------------------------|--------------------------------------|-----------------|
| | | Preventative* | Curative** |
| Dollar spot | 7-14 days | 95-190 | 190-350 |
| Helminthosporium leafspot and melting out | 7-10 days | 95-190 | 190-350 |
| Large brown patch | 7-10 days | 95-190 | 390-350 |
| Anthracnose | 7-10 days | 170-190 | not-recommended |

| | | | |
|---|--|-------------------|------------------------|
| Gray snow mold (Typhula spp.) and Pink snow mold (Gerlachia or Fusarium patch) | | 240-500*** | not-recommended |
|---|--|-------------------|------------------------|

- * Recommended rates for preventing disease establishment; use lower rate as when disease conditions are light to moderate, higher indicated rates when conditions are severe.
- ** Rates for use on a 7 day schedule when disease is present. Higher indicated rate should be applied under severe conditions.
- *** Apply DACONIL 2787 FLOWABLE FUNGICIDE at 500 milliliters or apply 240 milliliters as a tank mix with 125 to 250 milliliters of ROVRAL® Green 240 g/L per 100 sq. metres prior to snow cover in the fall and repeat in the early spring if conditions for snow mold continue to exist.

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.