

**GROUP 22 HERBICIDE****GRAMOXONE®**LIQUID HERBICIDE WITH WETTING AGENT  
(Contains Paraquat)

AGRICULTURAL

A non-residual Herbicide for the Control of Many Grasses and Broadleaf Weeds.  
Inactivated on Contact with the Soil.**FOR USE ONLY BY FARMERS AND PEST CONTROL OPERATORS.  
NOT FOR USE BY HOME GARDENERS.****GUARANTEE:**Paraquat .....200 g per litre (present as dichloride)  
..... This product contains 0.006 % formaldehyde**KEEP OUT OF REACH OF CHILDREN****DANGER****POISON****DANGER - CORROSIVE TO EYES**CALL A DOCTOR OR CONTACT A POISON CONTROL CENTRE IMMEDIATELY  
IN CASE OF ACCIDENT**WARNING****READ THE LABEL AND BOOKLET BEFORE USING.**

Danger - May be fatal if swallowed.

Never transfer to other containers.

Keep out of reach of children and animals.

Store tightly closed in original container, and in a safe place.

REGISTRATION NO. **8661**  
PEST CONTROL PRODUCTS ACTNET CONTENTS: **1 L - 415 L****Syngenta Crop Protection Canada, Inc.**  
140 Research Lane, Research Park  
Guelph, ON N1G 4Z3  
Telephone: 1-877-964-3682

## NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

## FIRST AID

**IN CASE OF POISONING**, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**If swallowed**, induce vomiting if not already occurring. Get to nearest hospital **FAST**. **THIS IS ESSENTIAL**. If delay unavoidable, administer fluids and induce further vomiting. Contact a poison control centre **IMMEDIATELY**.

**If in eyes**, flush with clean water for 15 minutes and get medical attention **IMMEDIATELY**.

**If on skin**, wash thoroughly with water. Remove contaminated clothing **IMMEDIATELY**; wash before re-use.

## TOXICOLOGICAL INFORMATION

**If swallowed**, give stomach wash-out and test urine and gastric aspirate for paraquat. If positive, give up to 1 litre of adsorbent suspension (activated charcoal) mixed with a purgative ( $\text{MgSO}_4$ ,  $\text{Na}_2\text{SO}_4$  or mannitol). Repeat administration of adsorbent suspension for the next 24 hours, plus purgatives as required. Maintain and monitor electrolyte and fluid status daily. Consider haemodialysis or haemoperfusion, using charcoal column. Delay oxygen as long as possible.

**If in eyes**, treat symptomatically, using antibiotics and steroids as necessary.

## PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

**DO NOT USE GRAMOXONE LIQUID HERBICIDE WITH WETTING AGENT UNDILUTED.** The maximum spray concentration used should not exceed 2.5 parts GRAMOXONE LIQUID HERBICIDE with WETTING AGENT to 100 parts water. **DANGER** - Corrosive to eyes. **PREVENT** all contact of the concentrate with the skin and eyes. **WEAR LONG-SLEEVE CHEMICAL RESISTANT GLOVES** (gauntlet style), **SAFETY GOGGLES**, **FACE SHIELD**, **AND LONG-SLEEVED SHIRT AND LONG PANTS OR COVERALLS AND A CHEMICAL RESISTANT APRON** when handling the concentrate, during mixing/loading and during application via handheld equipment. **WEAR LONG-SLEEVED SHIRT AND LONG PANTS OR COVERALLS** during application. **WEAR COVERALLS OVER A LONG-SLEEVED SHIRT AND LONG PANTS DURING APPLICATION WITH A BACKPACK SPRAYER.** **WEAR LONG-SLEEVED CHEMICAL RESISTANT GLOVES, A LONG-SLEEVED SHIRT AND LONG PANTS** during clean-up and repair

activities. Do not apply with a mist blower or high-pressure sprayer producing fine droplets. **AVOID WORKING** in spray mist. **AVOID CONTACT** with spray solution. **REMOVE HEAVILY CONTAMINATED CLOTHING IMMEDIATELY** and wash before re-use. **WASH** hands and face thoroughly after spraying and before smoking or taking meals. **DO NOT CONTAMINATE FEEDS, FOODSTUFFS, or WATER SUPPLIES. STORE IN ORIGINAL CONTAINER** tightly closed, in a safe place away from children. Overspray or drift to sensitive habitats should be avoided. **DO NOT** reenter treated areas within 24 hours. If required, individuals may reenter treated areas within 24 hours for short-term tasks not involving hand labour if wearing a long-sleeved shirt and long pants provided at least 4 hours have passed since applications.

## ENVIRONMENTAL HAZARDS

**TOXIC** to aquatic organisms and terrestrial plants. Observe buffer zones specified under **DIRECTIONS FOR USE**.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at [www.croplife.ca](http://www.croplife.ca).

## STORAGE

**STORE ABOVE 0°C.** If crystallization occurs because of storage below 0°C, warm to room temperature, agitate until reconstituted.

## SPILL CLEANUP

Wear appropriate protective equipment (gloves, glasses, apron) when attempting to clean up the spill. If the container is leaking, secure leak and place the container into a drum or heavy gauge plastic bag. Contact Syngenta Crop Protection Canada, Inc. (See EMERGENCY TELEPHONE NUMBER) for further information.

For spills and leaks, contain the liquid with dikes of inert material (soil, clay, kitty litter etc.). Absorb the spill onto inert material and shovel into a sealable waste container.

On hard surfaces - sprinkle spill area with detergent and scrub in a small quantity of water with a coarse broom. Let stand 10 minutes then absorb onto an inert material and shovel into the waste container. Dispose of all waste including scrub brush in accordance with provincial requirements.

On soil - remove the top 15 cm of soil in the spill area and replace with fresh soil.

## DECONTAMINATION AND DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean up of spills.

## **CONTAINER DISPOSAL:**

### **For recyclable containers:**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

### **For refillable containers:**

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not use this container for any other purpose.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING  
CALL 1-800-327-8633 (FASTMED)***

Product names marked ® or ™ are registered trademarks of a Syngenta Group Company

**GROUP 22 HERBICIDE**

**GRAMOXONE®**

LIQUID HERBICIDE WITH WETTING AGENT  
(Contains Paraquat)

AGRICULTURAL

A non-residual Herbicide for the Control of Many Grasses and Broadleaf Weeds.  
Inactivated on Contact with the Soil.

**FOR USE ONLY BY FARMERS AND PEST CONTROL OPERATORS.  
NOT FOR USE BY HOME GARDENERS.**

**GUARANTEE:**

Paraquat .....200 g per litre (present as dichloride)  
..... This product contains 0.006 % formaldehyde

**KEEP OUT OF REACH OF CHILDREN**

**DANGER**



**POISON**

**DANGER - CORROSIVE TO EYES**

CALL A DOCTOR OR CONTACT A POISON CONTROL CENTRE IMMEDIATELY  
IN CASE OF ACCIDENT

**WARNING**

**READ THE LABEL AND BOOKLET BEFORE USING.**

Danger - May be fatal if swallowed.

Never transfer to other containers.

Keep out of reach of children and animals.

Store tightly closed in original container, and in a safe place.

REGISTRATION NO. **8661**  
PEST CONTROL PRODUCTS ACT

**Syngenta Crop Protection Canada, Inc.**  
140 Research Lane, Research Park  
Guelph, ON N1G 4Z3  
Telephone: 1-877-964-3682

## NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

## FIRST AID

**IN CASE OF POISONING**, contact a physician or a poison control centre **IMMEDIATELY**. Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**If swallowed**, induce vomiting if not already occurring. Get to nearest hospital **FAST**. **THIS IS ESSENTIAL**. If delay unavoidable, administer fluids and induce further vomiting. Contact a poison control centre **IMMEDIATELY**.

**If in eyes**, flush with clean water for 15 minutes and get medical attention **IMMEDIATELY**.

**If on skin**, wash thoroughly with water. Remove contaminated clothing **IMMEDIATELY**; wash before re-use.

## TOXICOLOGICAL INFORMATION

**If swallowed**, give stomach wash-out and test urine and gastric aspirate for paraquat. If positive, give up to 1 litre of adsorbent suspension (activated charcoal) mixed with a purgative ( $\text{MgSO}_4$ ,  $\text{Na}_2\text{SO}_4$  or mannitol). Repeat administration of adsorbent suspension for the next 24 hours, plus purgatives as required. Maintain and monitor electrolyte and fluid status daily. Consider haemodialysis or haemoperfusion, using charcoal column. Delay oxygen as long as possible.

**If in eyes**, treat symptomatically, using antibiotics and steroids as necessary.

## PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

**DO NOT USE GRAMOXONE LIQUID HERBICIDE WITH WETTING AGENT UNDILUTED.** The maximum spray concentration used should not exceed 2.5 parts GRAMOXONE LIQUID HERBICIDE with WETTING AGENT to 100 parts water. **DANGER** - Corrosive to eyes. **PREVENT** all contact of the concentrate with the skin and eyes. **WEAR LONG-SLEEVE CHEMICAL RESISTANT GLOVES** (gauntlet style), **SAFETY GOGGLES**, **FACE SHIELD**, **AND LONG-SLEEVED SHIRT AND LONG PANTS OR COVERALLS AND A CHEMICAL RESISTANT APRON** when handling the concentrate, during mixing/loading and during application via handheld equipment. **WEAR LONG-SLEEVED SHIRT AND LONG PANTS OR COVERALLS** during application. **WEAR COVERALLS OVER A LONG-SLEEVED SHIRT AND LONG PANTS DURING APPLICATION WITH A BACKPACK SPRAYER.** **WEAR LONG-SLEEVED CHEMICAL RESISTANT GLOVES, A LONG-SLEEVED SHIRT AND LONG PANTS** during clean-up and repair

activities. Do not apply with a mist blower or high-pressure sprayer producing fine droplets. **AVOID WORKING** in spray mist. **AVOID CONTACT** with spray solution. **REMOVE HEAVILY CONTAMINATED CLOTHING IMMEDIATELY** and wash before re-use. **WASH** hands and face thoroughly after spraying and before smoking or taking meals. **DO NOT CONTAMINATE FEEDS, FOODSTUFFS, or WATER SUPPLIES. STORE IN ORIGINAL CONTAINER** tightly closed, in a safe place away from children. Overspray or drift to sensitive habitats should be avoided. **DO NOT** reenter treated areas within 24 hours. If required, individuals may reenter treated areas within 24 hours for short-term tasks not involving hand labour if wearing a long-sleeved shirt and long pants provided at least 4 hours have passed since applications.

## ENVIRONMENTAL HAZARDS

**TOXIC** to aquatic organisms and terrestrial plants. Observe buffer zones specified under **DIRECTIONS FOR USE**.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at [www.croplife.ca](http://www.croplife.ca).

## STORAGE

**STORE ABOVE 0°C.** If crystallization occurs because of storage below 0°C, warm to room temperature, agitate until reconstituted.

## SPILL CLEANUP

Wear appropriate protective equipment (gloves, glasses, apron) when attempting to clean up the spill. If the container is leaking, secure leak and place the container into a drum or heavy gauge plastic bag. Contact Syngenta Crop Protection Canada, Inc. (See EMERGENCY TELEPHONE NUMBER) for further information.

For spills and leaks, contain the liquid with dikes of inert material (soil, clay, kitty litter etc.). Absorb the spill onto inert material and shovel into a sealable waste container.

On hard surfaces - sprinkle spill area with detergent and scrub in a small quantity of water with a coarse broom. Let stand 10 minutes then absorb onto an inert material and shovel into the waste container. Dispose of all waste including scrub brush in accordance with provincial requirements.

On soil - remove the top 15 cm of soil in the spill area and replace with fresh soil.

## DECONTAMINATION AND DISPOSAL

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean up of spills.

## CONTAINER DISPOSAL:

### **For recyclable containers:**

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

### **For refillable containers:**

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not use this container for any other purpose.

***IN CASE OF EMERGENCY INVOLVING A MAJOR SPILL, FIRE OR POISONING  
CALL 1-800-327-8633 (FASTMED)***

## PRODUCT INFORMATION

GRAMOXONE LIQUID HERBICIDE with WETTING AGENT is a non-volatile, fast acting herbicide. It is inactivated on contact with the soil, therefore, has no residual effect in the soil. The herbicidal effect varies with weed species; hence, repeat applications may be necessary upon certain perennial weeds. Annual weeds are generally killed with one application if the weed growth has been completely covered with the spray solution.

GRAMOXONE LIQUID HERBICIDE with WETTING AGENT works rapidly, and rain falling shortly after application normally will not reduce the effectiveness of the treatment. APPLICATIONS MADE ON CLOUDY DAYS, DURING DULL SUNLIGHT OR JUST PRIOR TO OR DURING PERIODS OF DARKNESS WILL GENERALLY INCREASE THE SUBSEQUENT EFFECTIVENESS OF THE TREATMENT.

Thoroughly wet all foliage. For dense weed growth, use the higher rate and the higher volume of water.

**NOTE:** Do not add AGRAL<sup>®</sup> 90 as this GRAMOXONE LIQUID HERBICIDE with WETTING AGENT formulation already contains a wetting agent.

**CAUTIONS:** Avoid application or drift onto crops, ornamental plants, lawns, grazing areas or other desirable growth. Use clean (non-turbid) water for spraying GRAMOXONE LIQUID HERBICIDE with WETTING AGENT. Muddy water will reduce the effectiveness of the chemical. DO NOT APPLY THROUGH MIST BLOWERS. It is important to thoroughly wash equipment after spraying - Use a wetting agent (AGRAL 90 at 60 mL per 100 L of water), flush and spray out, then thoroughly rinse with clean water. When possible, the equipment should be filled with clean water and left overnight. Spray out before storing equipment or using for other materials.



DO NOT APPLY BY AIR.

## **DIRECTIONS FOR USE:**

### **GRASS AND WEED CONTROL IN FRUIT CROPS AND SHELTERBELTS:**

Rate and Method of Application: Apply 5.5 L GRAMOXONE LIQUID HERBICIDE with WETTING AGENT in 1100 L of water per sprayed hectare or 75 mL in 10 L of water per 100 m<sup>2</sup>. Of this mixture, 550 mL will treat an area 1.75 m in diameter around a tree. Applications of this product in fruit crops and shelterbelts must be made using low boom sprayers fitted with drift-reducing shrouds or shields. Follow manufacturer's recommendations for use of shrouds or shields with particular attention to maintaining the minimum allowable boom height. Use flat fan nozzles with the highest flow rate and lowest pressure that will provide good coverage, within the manufacturer's recommended range.

**APPLES AND GRAPES ONLY** - For hard-to-kill annual and perennial weeds, add 2.75 L (or 40 mL per 100 m<sup>2</sup>) of TROPOTOX<sup>†</sup> Plus 64 or TROPOTOX<sup>†</sup> to the GRAMOXONE LIQUID HERBICIDE with WETTING AGENT spray. Apply as directed for GRAMOXONE LIQUID HERBICIDE with WETTING AGENT except for field bindweed control under grape vines, apply when the field bindweed begins to bloom. Apply as directed spray to the base of trees, bushes, canes, and vines. On all crops listed below, avoid contact with foliage (especially on conifers), fruit, young growing shoots, suckers and green bark.

**UNDER APPLES, APRICOTS, CHERRIES, RED CURRANTS, GOOSEBERRIES, Highbush BLUEBERRIES, PEACHES, PEARS AND PLUMS** - Established one or more years: Repeat application as necessary.

**UNDER GRAPE VINES** - Established on trellises - Apply in May or early June and repeat if necessary. An area 1 m by 10,000 m of row equals one hectare.

**UNDER BLACKBERRIES, LOGANBERRIES AND RED RASPBERRIES** - Apply early in spring before new shoots emerge or in fall when canes are mature.

**UNDER ESTABLISHED SHELTERBELTS** - Repeat application if necessary.

For residual control of germinating annual weeds under Apples, Pears, Loganberries, Red Raspberries, Highbush Blueberries, Blackberries and Shelterbelts, GRAMOXONE LIQUID HERBICIDE with WETTING AGENT can be tank mixed with PRINCEP<sup>®</sup> NINE-T<sup>®</sup>. Use the appropriate rates for each crop and observe the precautions and limitations listed on the PRINCEP NINE-T label.

### **FILBERTS and HAZELNUTS: GREEN SUCKERS:**

As a directed spot treatment at the base of each tree, apply 75 mL of GRAMOXONE LIQUID HERBICIDE with WETTING AGENT in 10 L of water on green suckers to runoff.

For inter-row general weed and green sucker control performed in one operation using a boom sprayer, apply 2.8 to 5.6 L of GRAMOXONE LIQUID HERBICIDE with WETTING AGENT in 550 to 1100 L of water per hectare.

For residual control of germinating annual weeds, apply GRAMOXONE LIQUID HERBICIDE with WETTING AGENT at 5.0 L per hectare tank mixed with PRINCEP NINE-T, at 2.0 to 2.5 kg per hectare. Apply under trees established one year or more. Apply only one (1) application per year, by

ground. Consult the PRINCEP NINE-T label for additional information regarding choice of rate (as determined by soil type), precautions, restrictions and directions for mixing and application.

#### **INTER-ROW SPRAYING IN STRAWBERRIES TO CONTROL EXCESS PLANTS AND WEEDS:**

To obtain narrow strawberry rows in the planting year, or from just after harvest to fall in renovated plantings, apply 5.5 L of GRAMOXONE LIQUID HERBICIDE with WETTING AGENT in 550 to 1100 L of water per hectare to give thorough coverage of foliage. Determine width of the strawberry row to be retained and use shields to keep spray off the desired strawberry plants. Thus, the spray will be applied only to the area between the rows killing excess strawberry plants and weeds. In renovated plantings, 2 applications may be required - one just after harvest and one again in the fall. Apply on a calm day and use a spray nozzle arrangement and low pressure to avoid spray drift around the shields.

#### **WEED CONTROL BY STALE SEED BED TECHNIQUE FOR VEGETABLES AND FIELD CROPS:**

Pre-emergent to crop. Post emergent to weeds: For weed control in beans (all types) beets, carrots, cole crops, corn, cucumbers, onions, peas, potatoes, soybeans, and turnips, prepare a seed bed at least 2 to 4 weeks before seeding to stimulate weed growth. Seed without further cultivation and with a minimum disturbance of soil. To burn off emerged weeds apply 2.75 to 5.5 L GRAMOXONE LIQUID HERBICIDE with WETTING AGENT in 300 to 1100 L of water per hectare prior to or after seeding. Do not apply later than 3 days before crop emergence. Use 5.5 L rate when weeds are above 5 cm in height and higher volume of water on dense weed growth.

#### **INTER-ROW DIRECTED CHEMICAL WEEDING OF VEGETABLE, FIELD AND ESTABLISHED NURSERY CROPS:**

For weed and grass control between the rows after weed emergence, use equipment and nozzles designed to prevent spray contact with the green foliage or other green parts of plants. Apply 2.75 to 5.5 L GRAMOXONE LIQUID HERBICIDE with WETTING AGENT in 300 to 550 L of water per hectare. Use 4.25 to 5.5 L when weeds are above 5 cm in height and higher volume of water on dense weed growth.

#### **ASPARAGUS:**

Apply a GRAMOXONE LIQUID HERBICIDE with WETTING AGENT spray to emerged broadleaf weeds and grasses prior to, or after the cutting season. Do not apply to emerged spears which are to be used for human consumption. Apply 2.75 to 5.5 L GRAMOXONE LIQUID HERBICIDE with WETTING AGENT in 300 to 1100 L of water per hectare. Use 5.5 L rate when weeds are above 5 cm in height and higher volume of water on dense weed growth. For residual control of germinating annual weeds, GRAMOXONE LIQUID HERBICIDE with WETTING AGENT may be tank mixed with PRINCEP NINE-T at 2.5 to 3.75 kg/ha either 7 days before first cutting and/or after last harvest. Observe all precautions on the PRINCEP NINE-T label. **Do not spray the asparagus fern.**

#### **WEED CONTROL IN POTATOES:**

Apply 2.75 to 4.25 L in 300 to 550 L of water per hectare to control emerged quackgrass, annual grasses and broadleaf weeds. To control emerged seedling annual grasses and seedling annual broadleaf weeds only, apply 1.5 L in 300 to 550 L of water per hectare. Grasses and weeds germinating after spraying will require cultivation for control. For all rates of use, apply to potato varieties as follows: Netted Gem (Idaho Baker, Russett or Burbank) and Cherokee apply up to

ground crack only (potato tops about to emerge). For other varieties, apply up to time first potato tops have reached 5 to 8 cm in height (this height is less than it often appears to the eye when observing an emerging field).

GRAMOXONE LIQUID HERBICIDE with WETTING AGENT - Tank mixed with residual herbicides - For residual control of annual broadleaf weeds and annual grasses tank mix with one of the following:\*

PRODUCT	RATE
LOROX <sup>†</sup> L	2.25 - 4.5 L/ha
PATORAN <sup>†</sup> 400 SC	2.5 - 7.0 L/ha
AFESIN <sup>†</sup> 2EC	5.5 - 8.0 L/ha
LEXONE <sup>†</sup> DF	0.83 - 1.2 kg/ha
LEXONE L	1.3 - 1.8 L/ha
SENCOR <sup>†</sup> 500 FL	1.1 - 1.75 L/ha

\* Refer to the label of the residual herbicide used for specific rates on your soil type.

Apply to emerged weeds and grasses up to and including ground crack using the lower rate of residual herbicide on light soils and the higher rate on heavier and high organic matter soils. **Do not apply these tank mixes after ground crack when tops have emerged as excessive injury may result.** If quackgrass is present, use 2.75 to 4.25 L of GRAMOXONE LIQUID HERBICIDE with WETTING AGENT per hectare with the residual herbicide.

**NOTE:** OBSERVE ALL PRECAUTIONS AND USE DIRECTIONS FOR THE RESIDUAL HERBICIDE USED.

**NOTE:** Application to exposed or emerged potato foliage will cause temporary injury and chlorosis. Do not apply to emerged potato foliage in the evening, or when potatoes are under moisture stress due to extremely dry soil conditions or to early potatoes. The use of poor or diseased seed and cut seed with one eye will make potatoes more susceptible to injury by post- emergence GRAMOXONE LIQUID HERBICIDE with WETTING AGENT sprays. Treatment with GRAMOXONE LIQUID HERBICIDE with WETTING AGENT alone will normally eliminate several cultivations, but has no residual action and will not control growth of weeds which may take place following the application of GRAMOXONE LIQUID HERBICIDE with WETTING AGENT.

#### **ESTABLISHED ALFALFA FOR DEHYDRATION, HAY AND FORAGE:**

Apply GRAMOXONE LIQUID HERBICIDE with WETTING AGENT spray immediately (not later than 5 days) after cutting established alfalfa to control or suppress broadleaf weeds and grasses (including quackgrass) and release the alfalfa. Do not use a residual herbicide in the season prior to GRAMOXONE LIQUID HERBICIDE with WETTING AGENT application. Apply 2.75 to 5.5 L/ha in 300 to 1000 L of water.

#### **ESTABLISHED BIRDSFOOT TREFOIL FOR SEED:**

For improved seed yields and control of wild carrot, quackgrass and other broadleaf weeds and grasses, spray GRAMOXONE LIQUID HERBICIDE with WETTING AGENT at 2.75 to 5.5 L/ha in 300 to 500 L of water when the trefoil is 8 to 16 cm high and actively growing.

#### **ESTABLISHED BIRDSFOOT TREFOIL FOR HAY AND FORAGE:**

Apply GRAMOXONE LIQUID HERBICIDE with WETTING AGENT at 2.75 to 5.5 L/ha in 300 to 500

L of water immediately following the first cutting (within 5 days) in early June to control or suppress broadleaf weeds and grasses (including quackgrass) and release the legume.

#### **WEED CONTROL IN ROUGHLAND PASTURE RENOVATION WITH BIRDSFOOT TREFOIL:**

Apply a 2,4-D treatment in August to control perennial weeds prior to seeding with birdsfoot trefoil. Prior to full germination of birdsfoot trefoil in the spring, apply 5.5 L of GRAMOXONE LIQUID HERBICIDE with WETTING AGENT in 300 to 550 L of water per hectare to control grasses and annual weeds. This spray should be applied when the grasses are 5 to 10 cm high (early to mid-May) to give adequate suppression/control to allow establishment of the trefoil. Rotation grazing should be practiced in the establishment year to lessen injury to young seedlings.

#### **PASTURE RENOVATION - MINIMUM TILLAGE:**

Apply 2,4-D or a 2,4-D based treatment in August prior to treatment to control perennial broadleaf weeds. In May, after 5 to 10 cm of new growth has appeared apply 2.75 to 5.5 L of GRAMOXONE LIQUID HERBICIDE with WETTING AGENT in 300 to 500 L water per hectare to control annual weeds and grasses immediately prior to seeding trefoil or alfalfa with suitable sod seeding/minimum tillage equipment.

#### **PRE- OR POST-SEEDING BURNDOWN FOR BARLEY, CANARY SEED, CANOLA, CORN (field, sweet and pop types, including crops grown for seed production), DRY COMMON BEANS, FIELD (dry) PEAS, FLAX (including low linolenic acid varieties), LENTILS, MUSTARD, OATS, POTATOES, RYE, SOYBEANS, SUNFLOWERS, TRITICALE, WHEAT:**

For control of emerged annual grass and broadleaf weed seedlings, and top-growth control of perennial grass and broadleaf weeds apply 2.0 to 4.0 L per hectare. Apply in at least 100 L of water per hectare by ground sprayer. Apply before or after seeding, but prior to crop emergence to control emerged weeds. For spring control of winter annual weeds, or when weed growth is dense and weeds are greater than 10 cm in height, use at least 3.0 L per hectare and higher volumes of water. Do not graze or harvest crops within 30 days of treatment.

#### **CONSERVATION TILLAGE CORN:**

GRAMOXONE LIQUID HERBICIDE with WETTING AGENT may be applied in tank mixture with preemergence corn herbicides to control emerged annual grass and broadleaf weed seedlings and top growth of perennial grass and broadleaf weeds, and to provide residual control of weeds as claimed on the labels of the preemergence herbicides. Apply 2.75 to 4.0 L of GRAMOXONE LIQUID HERBICIDE with WETTING AGENT per hectare in 200 to 250 L water per hectare in combination with recommended rates of the following herbicides:

Product	Rate	Comments
AATREX <sup>®</sup> NINE-0 <sup>®</sup>	1.1 to 1.7 kg/ha	Use higher rate in fine textured soils and heavier weed populations
DUAL <sup>®</sup> MAGNUM <sup>®</sup> OR DUAL II MAGNUM	1.25 to 1.75 L/ha	Use higher rate for heavier weed populations
DUAL MAGNUM or DUAL II MAGNUM plus AATREX NINE-O	1.25 to 1.75 L/ha plus 1.1 to 1.7 kg/ha	Use higher rate of atrazine in fine textured soils
FRONTIER <sup>†</sup>	1.1 to 1.4 L/ha	Use the higher rate of atrazine on fine-textured soil
FRONTIER plus AATREX NINE-O	1.1 to 1.4 L/ha plus 1.1 to 1.7 kg/ha	Use the higher rate of atrazine and Frontier on fine-textured soil
PRIMEXTRA <sup>®</sup> II MAGNUM	3.0 – 4.0 L/ha	Use higher rate for heavier weed populations

Read and follow all precautions, restrictions, and directions on the labels of the tank mix partner herbicides.

### CONSERVATION TILLAGE SOYBEANS:

GRAMOXONE LIQUID HERBICIDE with WETTING AGENT may be applied in tank mixture with preemergence soybean herbicides to control emerged annual grass and broadleaf weed seedlings and top growth of perennial grass and broadleaf weeds, and to provide residual control of weeds as claimed on the labels of the preemergence herbicides. Apply 2.5 L of GRAMOXONE LIQUID HERBICIDE with WETTING AGENT per hectare in 200 to 250 L water per hectare in combination with recommended rates of the following herbicides:

Product	Rate	Comments
DUAL MAGNUM or DUAL II MAGNUM	1.15 to 1.75 L/ha	Use higher rate for heavier weed populations
DUAL MAGNUM or DUAL II MAGNUM plus SENCOR 500	1.15 to 1.75 L/ha plus 1.1 to 2.25 L/ha	Use higher rate for heavier weed populations
DUAL MAGNUM or DUAL II MAGNUM plus SENCOR 75 DF	1.15 to 1.75 L/ha plus 0.75 to 1.5 kg/ha	Use higher rate for heavier weed populations
DUAL MAGNUM or DUAL II MAGNUM plus SENCOR SOLUPAK	1.15 to 1.75 L/ha plus 0.75 to 1.5 kg/ha	Use higher rate for heavier weed populations
DUAL MAGNUM or DUAL II MAGNUM plus AFOLAN <sup>†</sup> F	1.15 to 1.75 L/ha plus 1.6 to 2.5 L/ha	Use higher rate for heavier weed populations
DUAL MAGNUM or DUAL II MAGNUM plus LOROX DF	1.15 to 1.75 L/ha plus 1.75 to 2.25 kg/ha	Use higher rate for heavier weed populations
DUAL MAGNUM or DUAL II MAGNUM plus LINURON 480	1.15 to 1.75 L/ha plus 1.9 to 2.3 L/ha	Use higher rate for heavier weed populations
BROADSTRIKE DUAL MAGNUM <sup>†</sup>	1.56 L/ha	
FRONTIER	1.1 to 1.4 L/ha	Use higher rate of Frontier on fine-textured or high organic soils
AFOLAN	2.7 to 5.0 L/ha	Use higher rate for muck and clay soils
LOROX DF	2.25 to 4.5 kg/ha	Use higher rate for muck and clay soils
LINURON <sup>†</sup> 480	2.25 to 4.5 L/ha	Use higher rate for muck and clay soils

Product	Rate	Comments
SENCOR 500 F	1.1 to 2.25 L/ha	Apply after planting but no later than 3 days prior to crop emergence. Apply when weeds are less than 4 cm in height.
SENCOR 75 DF	0.75 to 1.5 kg/ha	
SENCOR SOLUPAK	0.75 to 1.5 kg/ha	
PURSUIT <sup>†</sup>	0.312 to 0.42 L/ha	Apply preplant, up to 30 days before planting.

Read and follow all precautions, restrictions, and directions on the labels of the tank mix partner herbicides.

### USE PRECAUTIONS:

To reduce runoff from treated areas into aquatic habitats, consider the characteristics and conditions of the site before treatment. Site characteristics and conditions that may lead to runoff include, but are not limited to: heavy rainfall, moderate to steep slope, bare soil, poorly draining soil (e.g., soils that are compacted or fine textured such as clay). Avoid application of this product when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

**DO NOT** apply this product directly to aquatic areas.

**DO NOT** contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the ASAE medium classification.

### DO NOT APPLY BY AIR.

### BUFFER ZONES:

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, rangelands, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands).

Buffer zones are not required for use on fruit crops and shelterbelts, or for interrow directed chemical weeding of vegetable, field and established nursery crops as this product must be applied with low boom spray equipment fitted with drift-eliminating shrouds or shields specifically designed to prevent herbicide contact with sensitive vegetation.

Method of Application	Crop	Buffer Zones (metres) required for the protection of:			
		Freshwater habitats of depths			Terrestrial Habitats
		Less than 1 m	1-3 m	Greater than 3 m	
Field Sprayer*	Conservation Tillage Soybeans	40	30	20	15
	All Other Field Crops	50	40	30	20

\*Field sprayers fitted with drift-reducing devices can have their buffer zones reduced by 70% with the use of

shrouds or 30% with the use of cones.

When a tank mixture is used, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.

### **Resistance-Management Recommendations**

For resistance management, GRAMOXONE LIQUID HERBICIDE with WETTING AGENT is a Group 22 herbicide. Any weed population may contain or develop plants naturally resistant to GRAMOXONE LIQUID HERBICIDE with WETTING AGENT and other Group 22 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

#### **To delay herbicide resistance:**

Where possible, rotate the use of GRAMOXONE LIQUID HERBICIDE with WETTING AGENT or other Group 22 herbicides with different herbicide groups that control the same weeds in a field.

Use tank mixtures with herbicides from a different group when such use is permitted.

Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.

Monitor treated weed populations for resistance development.

Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.

Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.

For further information or to report suspected resistance, contact Syngenta Crop Protection Canada, Inc. company representatives at 1-87-SYNGENTA (1-877-964-3682) or at [www.syngenta.ca](http://www.syngenta.ca).

Product names marked ® or ™ are registered trademarks of a Syngenta Group Company

† All other products listed are registered trademarks of their respective companies.

\*\*\*\*\*

This label transcript service is offered by the Pest Management Regulatory Agency to provide efficient searching for label information. This service and this information do not replace the official hard-copy label. The PMRA does not provide any guarantee or assurance that the information obtained through this service is accurate, current or correct, and is therefore not liable for any loss resulting, directly or indirectly, from reliance upon this service.